

130 GeV gamma ray line and enhanced Higgs di-photon rate from Triplet-Singlet extended MSSM

Tanushree Basak* and Subhendra Mohanty†

Physical Research Laboratory, Ahmedabad 380009, India.

Abstract

We propose an economic extension of minimal supersymmetric standard model with a SU(2) singlet and $Y = 0$ triplet, which can explain (i) the 125 GeV Higgs boson without fine tuning, (ii) the 130 GeV γ -ray line seen at Fermi-LAT, (as well as a second photon line at 114 GeV)(iii) an enhanced Higgs di-photon decay rate seen by ATLAS, while being consistent with dark matter relic density and recent XENON 100 exclusion limits on spin-independent direct detection cross-section. We obtain the required cross-section of $10^{-27} \text{cm}^3 \text{s}^{-1}$ for the 130 GeV γ -ray flux through the resonant annihilation of dark matter via pseudoscalar triplet Higgs of mass ~ 260 GeV. The dark matter is predominantly bino-higgsino which has large couplings with photons (through higgsino) and gives correct relic density (through bino). We get the enhanced Higgs diphoton decay rate, $R_{\gamma\gamma} \simeq 1.24$ dominantly contributed by the light chargino-loops, which can account for the reported excess seen in the $h \rightarrow \gamma\gamma$ channel by ATLAS.

*Electronic address: tanu@prl.res.in

†Electronic address: mohanty@prl.res.in

1. INTRODUCTION

It is well-known that SUSY is the simplest model from protecting the Higgs mass from large radiative corrections without fine tuning. In the minimal supersymmetric standard model (MSSM) [1], the Higgs mass is close to the Z-boson mass at the tree level, which demands a large radiative correction to raise the Higgs mass to 125-126 GeV seen at the LHC [2, 3]. This in turn pushes the squark masses in the TeV range and hence the mixing in the top-stop sector becomes significant. This raises issues about fine tuning - which is somewhat solved by the so-called Next-to-minimal supersymmetric standard model (NMSSM) by adding a singlet chiral superfield to MSSM [4]. But to achieve a tree level Higgs mass close to 125 GeV, we need a large $\lambda SH_u.H_d$ coupling which borders in the nonperturbative regime of λ [5]. In [6], it was shown by adding a hypercharge $Y=0$ $SU(2)$ -triplet and a singlet chiral superfield there is an extra tree-level contribution to the Higgs mass and it can be raised close to 125 GeV at the tree level. Hence, no large contributions from stop loops is needed to get the required Higgs mass which alleviates the fine tuning problem of fixing the stop mass to a high precision at the GUT scale. In addition, the model contains a dark matter(DM) candidate of mass $\mathcal{O}(100)$ GeV, with a correct relic abundance.

Recently it has been pointed out [7–9] that the analysis of the Fermi-LAT gamma-ray data [10] reveals the existence of a peak at around 130 GeV coming from the vicinity of the galactic center. Further, it shows that the interpretation of the gamma ray peak as due to DM annihilation with mass $129.8 \pm 2.4^{+7}$ GeV and annihilation cross-section $\langle\sigma v\rangle_{\gamma\gamma} = (1.27 \pm 0.32_{-0.28}^{+0.18}) \times 10^{-27} cm^3 sec^{-1}$ fits the signal well. Numerous studies have been made to accommodate this feature in terms of DM annihilation in both model-independent way[11] and in specifically Standard Model (SM) extended by singlets and triplet [12–14]. After the discovery of the Higgs-like boson around mass window 125-126 GeV, there is another intriguing possibility of a signal beyond SM in the $h \rightarrow \gamma\gamma$ channel. The ratio between the Higgs di-photon decay rate observed at LHC and the one expected in the SM is $R_{\gamma\gamma} = 1.65_{-0.30}^{+0.34}$ for ATLAS ($m_h = 126$ GeV) and $R_{\gamma\gamma} = 0.78_{-0.26}^{+0.28}$ for CMS ($m_h = 125$ GeV) [15, 16]. This channel will be an important discriminator of models as future LHC data pinpoints this number. The implications of the modified diphoton decay width in a generic model independent approach have been discussed in ref.[17].

In MSSM, the neutralino LSP, being the favourite candidate for DM, annihilates into

two photons via loop-suppressed processes[18, 19] - the cross-section for which is usually too small to explain the signal. But, with a bino-like LSP [20] and through the exchange of light slepton and sneutrino the observed $\sigma v_{\gamma\gamma}$ is achieved in MSSM. An alternate possibility is to incorporate the internal Bremsstrahlung (IB), which can also give sharp spectral features in the γ -ray spectrum[21]. In bino DM annihilation to final state fermions, the fermion mass suppression in the cross section is avoided if there is a final state photon with the fermion pair [18, 19]. In ref.[22] it was pointed out that a significant higgsino component in the DM would lead to a continuum gamma ray spectrum from $W^\pm$ final states and would not be able to explain the gamma ray peak. To avoid this, IB from bino dominated LSP's is more promising but there is a problem in getting a natural SUSY model with 130 GeV bino DM which gives the correct the relic abundance. MSSM could accommodate the enhancement in the di-photon decay rate with highly mixed light staus and large $\tan\beta$ [23].

In addressing the problem of explaining the 130 GeV gamma ray features, NMSSM models are most widely studied[24, 25]. In NMSSM, the neutralino DM(~ 130 GeV) annihilates into two photon via resonant channel through pseudoscalar singlet Higgs ($m_{A_s} \sim 260$ GeV) and light charged particle loops. NMSSM can also successfully account for the excess seen in the $h \rightarrow \gamma\gamma$ channels[26], in the case of strong singlet-doublet mixing, although the partial width of $h \rightarrow b\bar{b}$ is highly reduced in these models. In a generalised version of NMSSM model(GNMSSM)[27] simultaneously both the signals from Fermi and LHC has been explained in the same benchmark scenario.

Enhancement of diphoton decay width has been studied well in the triplet extended SUSY models[28–31], where the contributions from to the charginos and charged Higgs(triplet like, with large triplet coupling) are taken into account. But, so far no benchmark points have been found which at the same time provide a viable DM in triplet extended SUSY models.

In the present paper, we attempt to explain the 130 GeV gamma ray spectral feature in the triplet-singlet extended MSSM[6] through the resonant annihilation of neutralino LSP into photons via pseudoscalar triplet Higgs of mass $\sim 2m_{DM}$, which couples to the DM via the Yukawa term, $\lambda_2 T_0 \tilde{H}_u^0 \cdot \tilde{H}_d^0$. In addition, our model predicts a second photon peak at around 114 GeV with the cross-section being 0.75 times $\langle\sigma v\rangle_{\gamma\gamma}$. This DM has a correct relic abundance of 0.109 where dominant contribution comes from $\langle\sigma v\rangle_{W+W^-}$. The spin-independent direct detection cross-section is well-below the latest XENON100[32] exclusion limits. Another motivation of this work is to provide an enhanced diphoton decay rate

compared to SM through the additional contribution from the light chargino loops. This would be a specific prediction of our model and can be tested in the future collider search.

This article is organised as follows: In section II, the model is described briefly mentioning the details about the superpotential, bound on the lightest Higgs mass and the fermionic sector. In the next section, we attempt to provide an explanation for the Fermi-LAT monochromatic gamma ray line features with a neutralino LSP pair annihilation into two photon via pseudoscalar Higgs triplet near resonance. We substantiate our claim with a specific benchmark scenario which satisfy all desired phenomenological requirements. Section III, shows a detail formulation of the diphoton Higgs decay width. We present a short summary and conclusions in the last section.

2. THE MODEL

By taking naturalness of the Higgs mass as a guiding criterion, we extend the superpotential of MSSM [6] by adding a SU(2) singlet and triplet chiral superfield S and T_0 respectively, where T_0 has hypercharge $Y = 0$,

$$\hat{T}_0 = \begin{pmatrix} \frac{\hat{T}_0^0}{\sqrt{2}} & -\hat{T}_0^+ \\ \hat{T}_0^- & \frac{-\hat{T}_0^0}{\sqrt{2}} \end{pmatrix} \quad (1)$$

The most general form of the superpotential can be written as,

$$\mathcal{W} = (\mu + \lambda \hat{S}) \hat{H}_d \cdot \hat{H}_u + \frac{\lambda_1}{3} \hat{S}^3 + \lambda_2 \hat{H}_d \cdot \hat{T}_0 \hat{H}_u + \lambda_3 \hat{S}^2 Tr(\hat{T}_0) + \lambda_4 \hat{S} Tr(\hat{T}_0 \hat{T}_0) + W_{Yuk}. \quad (2)$$

where the Yukawa part is same as in the MSSM. To solve the μ -problem, we reduce the general superpotential to the scale-invariant form as, which then possess an accidental Z_3 -symmetry,

$$W_{sc.inv.} = \lambda \hat{S} \hat{H}_d \cdot \hat{H}_u + \frac{\lambda_1}{3} \hat{S}^3 + \lambda_2 \hat{H}_d \cdot \hat{T}_0 \hat{H}_u + \lambda_4 \hat{S} Tr(\hat{T}_0 \hat{T}_0) + W_{Yuk}. \quad (3)$$

Therefore, an effective μ -term is generated when the neutral components of S and T_0 acquire vacuum expectation value (vev) v_s and v_t respectively,

$$\mu_{eff} = \lambda v_s - \frac{\lambda_2}{\sqrt{2}} v_t \quad (4)$$

Here, $v_u^2 + v_d^2 = v^2 = (174)^2 GeV^2$ (where, $\langle H_u^0 \rangle = v_u$, $\langle H_d^0 \rangle = v_d$) and $\tan \beta = \frac{v_u}{v_d}$.

Due to the addition of triplet, the ρ -parameter deviates from unity by a factor of $4 \frac{v_t^2}{v^2}$ at the tree level. It poses strong constraint on the triplet vev v_t from the Electroweak (EW) precision tests such that, $v_t \leq 9$ GeV [33] at 95% C.L.

The scalar potential of this model consists of three parts,

$$V = V_{SB} + V_F + V_D \quad (5)$$

where, V_{SB} consists of the soft-supersymmetry breaking term associated with the superpotential in equation(3),

$$V_{SB} = m_{H_u}^2 [|H_u^0|^2 + |H_u^+|^2] + m_{H_d}^2 [|H_d^0|^2 + |H_d^-|^2] + m_S^2 |S|^2 + m_T^2 \text{Tr}(T_0^\dagger T_0) + (-\lambda A_\lambda S H_u \cdot H_d + \frac{\lambda_1}{3} A_{\lambda_1} S^3 + \lambda_2 A_{\lambda_2} H_d \cdot T_0 H_u + \lambda_4 B_\lambda S \text{Tr}(T_0^2) + h.c) \quad (6)$$

V_F and V_D are the supersymmetric potential derived from F-terms and D-terms[6] respectively.

The CP-even higgs sector consists of four massive higgs as h , H_1 , H_2 and H_3 . Scalar parts of the singlet and triplet contribute significantly in the enhancement of the bound[34] on the lightest physical Higgs mass at the tree level as,

$$m_h^2 \leq M_Z^2 \left[\cos^2 2\beta + \frac{2\lambda^2}{g_1^2 + g_2^2} \sin^2 2\beta + \frac{\lambda_2^2}{g_1^2 + g_2^2} \sin^2 2\beta \right] \quad (7)$$

For moderate values of λ and λ_2 , the tree level mass can be lifted so that no large radiative corrections from the stop sectors are required to obtain $m_h = 125 - 126$ GeV. This indeed reduces the fine-tuning for the EW scale compared to MSSM, NMSSM and other triplet extended SUSY models.

The CP-odd higgs sector contains three pseudo-scalar Higgs A_1 , A_2 and A_3 . It always contains a Goldstone mode G^0 , which gives mass to Z-boson, and can be written as,

$$G^0 = \cos \beta H_{d_I}^0 - \sin \beta H_{u_I}^0$$

Likewise, there are three massive charged higgs $H_1^\pm$, $H_2^\pm$ and $H_3^\pm$ and the Goldstone mode $G^\pm$ gives mass to the W-bosons.

In the fermionic sector, the neutral component of the triplet and singlet i.e, $\tilde{T}^0$ and $\tilde{S}$ mix with the higgsinos and the gauginos. The neutralino mass matrix, in the gauge basis $(\tilde{B}, \tilde{W}^0, \tilde{H}_d^0, \tilde{H}_u^0, \tilde{S}, \tilde{T}^0)$ reads,

$$\mathcal{M}_{\tilde{G}} = \begin{pmatrix} M_1 & 0 & -c_\beta s_w M_Z & s_\beta s_w M_Z & 0 & 0 \\ 0 & M_2 & c_\beta c_w M_Z & -s_\beta c_w M_Z & 0 & 0 \\ -c_\beta s_w M_Z & c_\beta c_w M_Z & 0 & -\mu_{eff} & -\lambda v_u & \frac{\lambda_2}{\sqrt{2}} v_u \\ s_\beta s_w M_Z & -s_\beta c_w M_Z & -\mu_{eff} & 0 & -\lambda v_d & \frac{\lambda_2}{\sqrt{2}} v_d \\ 0 & 0 & -\lambda v_u & -\lambda v_d & 2\lambda_1 v_s & 2\lambda_4 v_t \\ 0 & 0 & \frac{\lambda_2}{\sqrt{2}} v_u & \frac{\lambda_2}{\sqrt{2}} v_d & 2\lambda_4 v_t & 2\lambda_4 v_s \end{pmatrix} \quad (8)$$

where, M_1 , M_2 are the soft breaking mass for Bino and Wino respectively. The lightest neutralino $\tilde{\chi}_1^0$, being the lightest supersymmetric particle (LSP), turns out to be a viable dark matter(DM) candidate.

Similarly, the charged component of the triplet, $\tilde{T}^+$ and $\tilde{T}^-$ contribute to the chargino mass matrix. The chargino matrix in the gauge basis $\tilde{G}^+$ and $\tilde{G}^-$ is given by,

$$\mathcal{M}_{ch} = \begin{pmatrix} M_2 & \frac{1}{\sqrt{2}}g_2v_d & g_2v_t \\ \frac{1}{\sqrt{2}}g_2v_u & \lambda v_s + \frac{\lambda_2}{\sqrt{2}}v_t & \lambda_2v_d \\ -g_2v_t & \lambda_2v_u & 2\lambda_4v_s \end{pmatrix} \quad (9)$$

where,

$$\tilde{G}^+ = \begin{pmatrix} \tilde{W}^+ \\ \tilde{H}_u^+ \\ \tilde{T}^+ \end{pmatrix}, \quad \tilde{G}^- = \begin{pmatrix} \tilde{W}^- \\ \tilde{H}_d^- \\ \tilde{T}^- \end{pmatrix}$$

Since, $\mathcal{M}_{ch}^T \neq \mathcal{M}_{ch}$, this matrix is diagonalised via bi-unitary transformation, which requires two distinct unitary matrices U and V such that,

$$\begin{aligned} \tilde{\chi}^+ &= V\tilde{G}^+, \\ \tilde{\chi}^- &= U\tilde{G}^- \end{aligned} \quad (10)$$

The diagonal matrix reads,

$$U^*\mathcal{M}_{ch}V^{-1} = \begin{pmatrix} m_{\tilde{\chi}_1^\pm} & 0 & 0 \\ 0 & m_{\tilde{\chi}_2^\pm} & 0 \\ 0 & 0 & m_{\tilde{\chi}_3^\pm} \end{pmatrix} \quad (11)$$

and similarly the hermitian conjugate of eqn.11 also gives diagonal chargino mass matrix.

3. 130 GEV FERMI GAMMA RAY LINE

In this model, the dark matter is the LSP $\tilde{\chi}_1^0$ which can be expressed in the gauge basis as,

$$\tilde{\chi}_1^0 = N_{11}\tilde{B} + N_{12}\tilde{W}_3^0 + N_{13}\tilde{H}_d^0 + N_{14}\tilde{H}_u^0 + N_{15}\tilde{S} + N_{16}\tilde{T}^0 \quad (12)$$

where, N_{11}^2 is the bino-fraction, N_{12}^2 is the wino-fraction, $N_{13}^2 + N_{14}^2$ is the higgsino-fraction, N_{15}^2 and N_{16}^2 are the singlino and triplino-fraction respectively.




FIG. 1: The dominant diagram for the resonant pair annihilation of neutralino into two photons via pseudoscalar triplet Higgs A_T

We scan the corresponding regions of the parameter space of the triplet-singlet model [6] and tune the couplings and masses, such that they satisfy all desired phenomenological properties. In Table.I, we show a sample set of benchmark points for a particular choice of $\tan \beta = 1.8$ specifying all the parameters, couplings and soft masses at the EW scale.

- As shown in [6], the CP-even physical Higgs boson receives significant contribution from the singlet and triplet through the terms $\lambda \hat{S} \hat{H}_d \cdot \hat{H}_u$ and $\lambda_2 \hat{H}_d \cdot \hat{T}_0 \hat{H}_u$ and thus its mass is raised to 122.9 GeV at tree level. It requires a little contribution from the radiative corrections raise it to 126 GeV. This lightest CP-even Higgs is SM-like with large H_u^0 and H_d^0 component.
- A dominantly triplet-like pseudoscalar Higgs A_T with mass ~ 260.54 GeV can be obtained by adjusting the soft-trilinear couplings. The pseudoscalar triplet A_T has no tree-level coupling with the SM fermions or gauge bosons. It can only interact with the neutralinos and charginos via the Yukawa term in the lagrangian like $\lambda_2 A_T \tilde{H}_u^0 \cdot \tilde{H}_d^0$. Therefore the width of A_T is small, i.e, $\Gamma_T \simeq 1.59$ keV- which boosts the Breit-Wigner propagator and cross-section $\langle \sigma v \rangle_{\gamma\gamma}$.
- The LSP $\tilde{\chi}_1^0$ is dominantly bino-like ($N_{11} \sim 0.84$) but contains substantial higgsino-fraction ($N_{13} \sim -0.31$ and $N_{14} \sim 0.36$). By suitably tuning the soft masses M_1 and M_2 , the desired mass of 130 GeV is obtained. Varying M_1 between 150-160 GeV, we obtain $127 \leq M_{\tilde{\chi}^0} [\text{GeV}] \leq 133$. Here, $\mu\text{-eff} \sim 246$ GeV being less than $v_s \sim 450$ GeV makes the singlino ($N_{15} \sim -0.19$) and triplino-fraction ($N_{16} \sim 0.10$) less in $\tilde{\chi}_1^0$. Again, since M_1 is lighter

Parameters at EW scale	
$\tan \beta$	1.8
λ	0.55
λ_1	0.20
λ_2	0.80
λ_4	0.25
$\mu_{eff}[\text{GeV}]$	246
$A_\lambda[\text{GeV}]$	400
$A_{\lambda_1}[\text{GeV}]$	-50
$A_{\lambda_2}[\text{GeV}]$	297.6
$B_\lambda[\text{GeV}]$	270
$v_t[\text{GeV}]$	2
$M_1[\text{GeV}]$	154.5
$M_2[\text{GeV}]$	375
Higgs Spectrum [GeV]	
m_h^{Tree}	122.93
m_{H_1}	175.29
m_{H_2}	457.27
m_{H_3}	538.86
m_{A_1}	142.12
m_{A_2}	260.54
m_{A_3}	534.56
$m_{H_1}^\pm$	133.13
$m_{H_2}^\pm$	365.61
$m_{H_3}^\pm$	545.59

Neutralino Masses [GeV]	
$m_{\tilde{\chi}_1^0}$	130.02
$m_{\tilde{\chi}_2^0}$	189.0
$m_{\tilde{\chi}_3^0}$	215.47
$m_{\tilde{\chi}_4^0}$	269.30
$m_{\tilde{\chi}_5^0}$	283.49
$m_{\tilde{\chi}_6^0}$	414.20
Chargino Masses [GeV]	
$m_{\tilde{\chi}_1^\pm}$	131.92
$m_{\tilde{\chi}_2^\pm}$	299.38
$m_{\tilde{\chi}_3^\pm}$	422.24
Observables	
Ωh^2	0.109
$\sigma(p)_{SI}[10^{-9}\text{pb}]$	0.681
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow\gamma\gamma) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	1.249
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow Z\gamma) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	0.94
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow WW) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	3.57
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow ZZ) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	0.62
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow b\bar{b}) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	0.045
$\langle\sigma v\rangle(\chi_1^0\chi_1^0\rightarrow\tau\bar{\tau}) [10^{-27}\text{cm}^3\text{ s}^{-1}]$	0.082
$R_{\gamma\gamma}$	1.24

TABLE I: A sample set of benchmark points for $\tan \beta = 1.8$ and $M_1 = 154.5$ GeV. The mass spectrum indicates all masses at the tree-level

than μ -eff, we get a enhancement in the bino fraction compared to higgsino. But, the significant higgsino fraction is required to get large value of $\langle\sigma v\rangle_{\gamma\gamma}$ through the resonant annihilation via psedoscalar Higgs A_T and the light chargino loops. In FIG.1 the resonant annihilation channel into two photon is shown. The lightest chargino $\tilde{\chi}_1^+$ and the DM are almost degenarate and is also dominantly higgsino-like.

The pair annihilation of $\tilde{\chi}^0$, with mass $129.8 \pm 2.4^{+7}$ GeV into two photon demands a cross-section of $\langle\sigma v\rangle_{\gamma\gamma} = (1.27 \pm 0.32_{-0.28}^{+0.18}) \times 10^{-27} cm^3 sec^{-1}$ in order to fit the Fermi-LAT signal[10].

A simplified form of the analytical expression of $\langle\sigma v\rangle_{\gamma\gamma}$ following [11] ,

$$\langle\sigma v\rangle_{\gamma\gamma} = \frac{\alpha^2 g_f^2 g_\chi^2}{256\pi^3} \frac{m_{\chi_1^+}^2}{[(4m_{DM}^2 - m_{A_T}^2)^2 + \Gamma_T^2 m_{A_T}^2]} \times [\arctan[(m_{\chi_1^+}^2 - m_{DM}^2)/m_{DM}^2]^{-1/2}]^2 \quad (13)$$

where, g_χ and g_f are the couplings of psedoscalar Higgs A_T with DM and the charged fermion in the loop respectively. Here, we take the assumption that only the lightest chargino, with mass 131.9 GeV contributes significantly. Upto a crude approximation, $g_\chi \sim \lambda_2 N_{13} N_{14}$ and $g_f \sim \lambda_2 U_{12} V_{12}$, where U and V are diagonalising matrix for the charginos. Finally, in the resonance limit of $m_{A_T} \sim 2m_{DM}$ and $m_{\chi^+} \rightarrow m_{DM}$, the pair annihilation cross-section becomes $\sim 1.249 \times 10^{-27} cm^3 s^{-1}$. However, the mass of the triplet-like CP-odd scalar Higgs has to lie accidentally close to 260 GeV to a precision ≤ 1.5 GeV. FIG.2 shows the behaviour of $\sigma v_{\gamma\gamma}$ with the psedoscalar triplet mass near resonance, this clarifies the need of tuning of both $M_{\tilde{\chi}_1^0}$ and m_{A_T} .

- **A second γ -ray line at 114 GeV :** Apart from the monochromatic γ -ray line at 130 GeV, there is another intriguing hint for a second line at ~ 111 GeV [35, 36], where the best fit to the relative cross-section is $\langle\sigma v\rangle_{\gamma Z}/\langle\sigma v\rangle_{\gamma\gamma} = 0.66_{-0.48}^{+0.71}$ [9]. A second photon line at 114 GeV is expected from kinematics if there is a $Z\gamma$ final state in the annihilation of $\tilde{\chi}_1^0$,

$$E_\gamma = m_{\tilde{\chi}_1^0} \left(1 - \frac{m_Z^2}{4m_{\tilde{\chi}_1^0}^2}\right) \quad (14)$$

where, $E_\gamma = 114$ GeV for $m_{\tilde{\chi}_1^0} = 130$ GeV. The cross-section for $\langle\sigma v\rangle_{\gamma Z}$ is calculated using an approximation of the formulae given in [37]. Here, we find that for the set of benchmark points presented in Table.I, $\langle\sigma v\rangle_{\gamma Z} \simeq 0.943 \times 10^{-27} cm^3 s^{-1}$.

Relic Density :

Another issue with dark matter is to satisfy the correct relic abundance, which is difficult in case




FIG. 2: Plot of $\sigma v_{\gamma\gamma}$ as a function of pseudoscalar mass M_{A_T} . The dashed line shows the maximum value of $\langle\sigma v\rangle_{\gamma\gamma} \simeq 1.249 \times 10^{-27} \text{cm}^3 \text{s}^{-1}$.

when it is dominantly higgsino-like since it couples to gauge boson very efficiently and thus leads to large pair annihilation cross-section. This kind of interaction can be reduced by an enhanced bino component. We find a neutralino DM with $N_{11} \sim 0.84$, $N_{13} \sim -0.31$ and $N_{14} \sim 0.36$, which makes the relic density 0.109. The pair annihilations into final state W^+W^- , ZZ , $b\bar{b}$, $\tau^+\tau^-$ are shown in Table.I, calculated using micrOMEGAs2.4 [38]. Thus, a bino dominated but with a substantial higgsino component dark matter is preferable in order to satisfy the latest PLANCK result, i.e, $\Omega_\chi h^2 = 0.1199 \pm 0.0027$ at 68% CL [39] whereas the corresponding value from the 9-year WMAP data is $\Omega_\chi h^2 = 0.1148 \pm 0.0019$ [40].

Calculation of spin-independent cross-section :

Starting from a low-energy neutralino-quark effective lagrangian for spin-independent interaction,

$$L_{eff} = a_q \bar{\tilde{\chi}}_1^0 \tilde{\chi}_1^0 \bar{q} q \quad (15)$$

where, a_q is the neutralino-quark coupling, we obtain the scattering cross section (spin-independent) for the dark matter off of a proton or neutron as,

$$\sigma_{scalar} = \frac{4m_r^2}{\pi} f_{p,n}^2 \quad (16)$$

where, m_r is the reduced mass of the nucleon and $f_{p,n}$ is the neutralino coupling to proton or

neutron[41, 42], given by

$$f_{p,n} = \sum_{q=u,d,s} f_{Tq}^{(p,n)} a_q \frac{m_{p,n}}{m_q} + \frac{2}{27} f_{TG}^{(p,n)} \sum_{q=c,b,t} a_q \frac{m_{p,n}}{m_q}, \quad (17)$$

where $f_{Tu}^{(p)} = 0.020 \pm 0.004$, $f_{Td}^{(p)} = 0.026 \pm 0.005$, $f_{Ts}^{(p)} = 0.118 \pm 0.062$, $f_{Tu}^{(n)} = 0.014 \pm 0.003$, $f_{Td}^{(n)} = 0.036 \pm 0.008$ and $f_{Ts}^{(n)} = 0.118 \pm 0.062$ [43]. $f_{TG}^{(p,n)}$ is related to these values by

$$f_{TG}^{(p,n)} = 1 - \sum_{q=u,d,s} f_{Tq}^{(p,n)}. \quad (18)$$

In deriving an approximate form of a_q/m_q we ignore contributions from the squark exchange diagrams because of the latest LHC bounds on squark masses[44, 45]. Thus, a_q receives significant contribution from the t-channel exchange of CP-even Higgs bosons. The analytical form of a_q goes roughly as,

$$\frac{a_q}{m_q} \simeq \frac{S_{\chi\chi h_i}}{m_{h_i}^2} S_{h_i q q} \quad (19)$$

where, $S_{\chi\chi h_i}$ is the coupling between the neutralino and the CP-even Higgs bosons. For, up-type quarks, $S_{h_i uu} = \frac{g_2}{2M_w \sin\beta} S_{i1}$ and down-type, $S_{h_i dd} = \frac{g_2}{2M_w \cos\beta} S_{i2}$. Now, the coupling $S_{\chi\chi h_i}$ is a product of different combinations of λ 's, N_{1k} and $S_{i1,2}$. S_{ij} is the matrix which diagonalises the CP-even Higgs matrix, and the weak eigenstate basis is $(H_{uR}^0, H_{dR}^0, T_R^0, S_R)$. N_{1k} 's are the different components of the lightest neutralino dark matter. Under the assumption that only the lightest physical Higgs boson, i.e, h_1 ($m_{h_1} \simeq 125.8$ GeV) contributes dominantly, $S_{\chi\chi h_1}$ takes the form,

$$\begin{aligned} S_{\chi\chi h_1} \simeq & g_2(N_{12} - \tan\theta_W N_{11})(S_{11}N_{13} - S_{12}N_{14}) \\ & - \sqrt{2}\lambda(S_{11}N_{14}N_{15} + S_{12}N_{13}N_{15} + S_{14}N_{14}N_{13}) + \sqrt{2}\lambda_1 S_{14}N_{15}^2 \\ & + \lambda_2(S_{11}N_{16}N_{13} + S_{12}N_{16}N_{14} + S_{13}N_{13}N_{14}) \\ & + \sqrt{2}\lambda_4(S_{14}N_{16}^2 + 2S_{13}N_{15}N_{16}) \end{aligned} \quad (20)$$

where the first term is the usual MSSM contribution, the second and third terms are due to the singlet. The fourth and fifth terms are the triplet contribution coming from $\lambda_2 H_d T_0 H_u$ and $\lambda_4 S Tr(T_0 T_0)$ in the superpotential respectively. Numerical values of the components S_{1j} ($j = 1, \dots, 4$) as obtained from the benchmark point are, $S_{11} \sim 0.885$, $S_{12} \sim 0.463$, $S_{13} \sim 0.026$ and $S_{14} \sim -0.037$. In this model, we find that the spin-independent cross-section $\sigma_p \simeq 6.8 \times 10^{-10}$ pb, which is well below the upper bound presented by the latest XENON 100 results[32] and can be accessible by the future XENON 1T experiment.

4. DI-PHOTON HIGGS DECAY RATE

In the SM, the diphoton decay of the Higgs boson is attributed through the W-boson loop and the contribution from the top-quark destructively interferes with the dominant W-boson contribution. The analytic expression for the diphoton partial width given as [46, 47]

$$\Gamma(h \rightarrow \gamma\gamma) = \frac{G_F \alpha^2 m_h^3}{128\sqrt{2}\pi^3} |A_1(\tau_W) + N_c Q_t^2 A_{1/2}(\tau_t)|^2, \quad (21)$$

where G_F is the Fermi constant, $N_c = 3$ is the number of color, $Q_t = +2/3$ is the top quark electric charge in units of $|e|$, and $\tau_i \equiv 4m_i^2/m_h^2$, $i = t, W$. The loop functions $A_1(\tau_W)$ and $A_{1/2}(\tau_t)$ for spin-1 (W boson) and spin-1/2 (top quark) particles are given in [48]. The numerical values of the loop functions for $m_h = 125$ GeV are,

$$A_1(\tau_W) \simeq -8.3, \quad A_{1/2}(\tau_t) \simeq 1.4$$

But in SUSY, we have additional contributions from the s-tops and charginos loops, which would significantly interfere with the SM contributions. Therefore, in general the branching width of Higgs decay to di-photon is formulated as [48],

$$\Gamma(h \rightarrow \gamma\gamma) = \frac{\alpha^2 m_h^3}{1024\pi^3} \left| \frac{g_{hVV}}{m_V^2} Q_V^2 A_1(\tau_V) + \frac{2g_{hff}}{m_f} N_{c,f} Q_f^2 A_{1/2}(\tau_f) + N_{c,S} Q_S^2 \frac{g_{hSS}}{m_S^2} A_0(\tau_S) \right|^2, \quad (22)$$

In the above the equation V , f , and S refer to generic spin-1, spin-1/2, and spin-0 particles, respectively. Q_V , Q_S and Q_f are the electric charges of the vectors, scalars and fermions in units of $|e|$, $N_{c,f}$ and $N_{c,S}$ are the number of fermion and scalar colors. $A_1(\tau_V)$, $A_{1/2}(\tau_f)$ and $A_0(\tau_S)$ are the loop functions for the vectors, fermions and scalars respectively.

In this model, the additional contribution to the diphoton Higgs decay width comes from the light chargino and the charged Higgs. Here, we take the assumption that the lightest charged Higgs (being dominantly triplet-like) only contribute to the decay width, since the other charged Higgs are much heavier. Now the term in the potential which gives rise to $hH^\pm H^\pm$ interaction is,

$$V_F \supset \lambda_2^2 v_u H_u^0 T_0^+ T_0^- \quad (23)$$

Therefore, the coupling $g_{hH^\pm H^\pm}$ becomes $\sim \lambda_2^2 v \sin \beta S_{11} C_{13} C_{14}$, where C_{ij} is the diagonalising matrix for the charged Higgs and $C_{13} \sim -0.669$, $C_{14} \sim -0.742$. The loop function for the scalar $A_0(\tau_s)$ is given by [48],

$$A_0(\tau_s) = -\tau_s^2 [\tau_s^{-1} - f(\tau_s^{-1})] \quad (24)$$




FIG. 3: Left Panel : Contours of $R_{\gamma\gamma}$ as a function of $\tan\beta$ and the triplet coupling λ_2 with $M_2 = 375$ GeV. Right Panel : Contours of $R_{\gamma\gamma}$ as a function of $\tan\beta$ and M_2 with $\lambda_2 = 0.8$

where, $f(\tau_s) = \arcsin^2 \sqrt{\tau_s}$ for, $\tau_s > 1$.

Therefore, considering the main contributions due to charginos, charged triplet, W -boson and top quark t and in the limit $m_h^2 \ll 4m_{\tilde{\chi}_i^\pm}^2$, the diphoton Higgs decay rate with respect to the SM value becomes[29],

$$R_{\gamma\gamma} = \left| 1 + \frac{\frac{4}{3} \frac{\partial}{\partial \log v} \log \det \mathcal{M}_{ch}(v) + \frac{g_{hH^\pm H^\pm}}{m_{H_1^\pm}^2} A_0(\tau_s)}{A_1(\tau_W) + \frac{4}{3} A_{1/2}(\tau_t)} \right|^2, \quad (25)$$

The numerator (first term) in Eq. (25) is given by

$$\frac{\partial}{\partial \log v} \log \det \mathcal{M}_{ch}(v) = - \frac{v^2 [\sin 2\beta (\lambda_2^2 M_2 + 2g_2^2 \lambda_4 v_s) - 2\lambda_2 g_2^2 v_t]}{2(M_2 \lambda_4 v_s + g_2^2 v_t^2) \mu_{eff} - \frac{1}{2} v^2 [\sin 2\beta (\lambda_2^2 M_2 + 2g_2^2 \lambda_4 v_s) - 2\lambda_2 g_2^2 v_t]}, \quad (26)$$

and its sign depends on the specific choices for the parameters. We are specifically interested in the region of parameter space where the numerator is negative (since the denominator is also negative), such that we obtain, $R_{\gamma\gamma} > 1$. We find that, the factor $g_{hH^\pm H^\pm}/m_{H_1^\pm}^2 \sim 0.0024$ and thus the contribution due to the extra charged triplet is treated to be negligible compared to the light chargino loops.

We see that for the set of benchmark points specified in Table.I, we obtain chargino masses in the range, $M_{\chi_i^\pm} \ni [131.92, 299.38, 422.24]$ GeV for $\tan\beta = 1.8$. This choice of parameter gives, $R_{\gamma\gamma} \simeq 1.224$. In FIG.3 (left panel), we show the contours of $R_{\gamma\gamma}$ in the $(\tan\beta, \lambda_2)$ plane for $M_2 = 375$ GeV. We observe that 50% enhancement is achieved with $\tan\beta \simeq 2$ but the triplet coupling $\lambda_2 (\geq 1.1)$ enters nonperturbative regime. Right panel of FIG.3 shows the dependence of $R_{\gamma\gamma}$ on $\tan\beta$ and M_2 . Here, we note that lowering the value of M_2 increases the $R_{\gamma\gamma}$, but then we deviate from other phenomenological requirements.

5. CONCLUSION

Recent analysis of the Fermi-LAT data shows existence of a monochromatic γ -ray line like features at $E_\gamma \sim 130$ GeV in the vicinity of the galactic center. A possible interpretation comes from DM annihilation into two photons, which demands the annihilation cross-section to be $1.27 \times 10^{-27} \text{cm}^3 \text{s}^{-1}$. A specific set of benchmark points is obtained which satisfy all desired phenomenological requirements including a 125 GeV lightest CP-even Higgs boson. In this work, we scan the parameter space of the triplet-singlet extended MSSM in order to obtain the required cross-section through the pair annihilation of 130 GeV neutralino DM via a pseudoscalar Higgs triplet of mass $M_{A_T} \sim 2m_{DM}$ near resonance and light chargino loops. The width of the pseudoscalar triplet being small helps in boosting the Breit-Wigner cross-section, $\langle\sigma v\rangle_{\gamma\gamma}$, since it only couples to the neutralinos and charginos via the Yukawa term $\lambda_2 A_T \tilde{H}_u^0 \tilde{H}_d^0$. Besides, this model predicts a second γ -ray peak at 114 GeV from the annihilation $\chi\chi \rightarrow \gamma Z$, and the cross section is approximately 0.75 times that of $\langle\sigma v\rangle_{\gamma\gamma}$, which is below the upper limit reported by Fermi LAT. The dark matter candidate being a mixture of bino-higgsino, leads to a relic abundance of 0.109, consistent with the PLANCK and WMAP-9 year data. The spin-independent scattering cross-section with nucleons is $0.68 \times 10^{-9} \text{pb}$, which is well below the latest XENON100 exclusion limits.

Latest data from ATLAS reveals an excess in diphoton decay width compared to SM, although CMS recent data is in tension with this claim. Our model predicts a similar enhancement in the diphoton decay rate as, $R_{\gamma\gamma} \sim 1.224$, which is contributed dominantly through the light chargino

loops. Such a prediction opens the possibility of this model being tested in future LHC runs.

- [1] H. Baer, V. Barger and A. Mustafayev, arXiv:1112.3017 [hep-ph];
S. Heinemeyer, O. Stal and G. Weiglein, Phys. Lett. B **710**, 201 (2012) [arXiv:1112.3026 [hep-ph]];
A. Arbey, M. Battaglia, A. Djouadi, F. Mahmoudi and J. Quevillon, Phys. Lett. B **708**, 162 (2012) [arXiv:1112.3028 [hep-ph]];
P. Draper, P. Meade, M. Reece and D. Shih, arXiv:1112.3068 [hep-ph];
A. Arbey, M. Battaglia and F. Mahmoudi, Eur. Phys. J. C **72**, 1906 (2012) [arXiv:1112.3032 [hep-ph]];
M. Kadastik, K. Kannike, A. Racioppi and M. Raidal, arXiv:1112.3647 [hep-ph];
J. Cao, Z. Heng, D. Li and J. M. Yang, arXiv:1112.4391 [hep-ph];
J. Ellis and K. A. Olive, arXiv:1202.3262 [hep-ph];
M. A. Ajaib, I. Gogoladze, F. Nasir and Q. Shafi, arXiv:1204.2856 [hep-ph];
F. Brummer, S. Kraml and S. Kulkarni, arXiv:1204.5977 [hep-ph].
- [2] G. Aad *et al.* [ATLAS Collaboration], “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” Phys. Lett. B **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]].
- [3] S. Chatrchyan *et al.* [CMS Collaboration], “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” Phys. Lett. B **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]].
- [4] M. Drees, Int. J. Mod. Phys. A **4**, 3635 (1989);
U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. **496**, 1 (2010);
G. G. Ross and K. Schmidt-Hoberg, arXiv:1108.1284 [hep-ph];
L. J. Hall, D. Pinner and J. T. Ruderman, arXiv:1112.2703 [hep-ph];
S. F. King, M. Muhlleitner and R. Nevzorov, Nucl. Phys. B **860**, 207 (2012) [arXiv:1201.2671 [hep-ph]];
Z. Kang, J. Li and T. Li, arXiv:1201.5305 [hep-ph];
J. Cao, Z. Heng, J. M. Yang, Y. Zhang and J. Zhu, JHEP **1203**, 086 (2012) [arXiv:1202.5821 [hep-ph]];

- D. A. Vasquez, G. Belanger, C. Boehm, J. Da Silva, P. Richardson and C. Wymant, arXiv:1203.3446 [hep-ph];
- U. Ellwanger and C. Hugonie, arXiv:1203.5048 [hep-ph].
- [5] S. F. King, M. Muhlleitner and R. Nevzorov, Nucl. Phys. B **860**, 207 (2012) [arXiv:1201.2671 [hep-ph]].
- [6] T. Basak and S. Mohanty, Phys. Rev. D **86**, 075031 (2012) [arXiv:1204.6592 [hep-ph]].
- [7] T. Bringmann, X. Huang, A. Ibarra, S. Vogl, C. Weniger and , JCAP **1207**, 054 (2012) [arXiv:1203.1312 [hep-ph]].
- [8] C. Weniger, JCAP **1208**, 007 (2012) [arXiv:1204.2797 [hep-ph]].
- [9] T. Bringmann and C. Weniger, Phys. Dark Univ. **1**, 194 (2012) [arXiv:1208.5481 [hep-ph]].
- [10] M. Ackermann *et al.* [LAT Collaboration], Phys. Rev. D **86**, 022002 (2012) [arXiv:1205.2739 [astro-ph.HE]].
- [11] M. R. Buckley and D. Hooper, Phys. Rev. D **86**, 043524 (2012) [arXiv:1205.6811 [hep-ph]].
- [12] J. M. Cline, Phys. Rev. D **86**, 015016 (2012) [arXiv:1205.2688 [hep-ph]].
- [13] B. Kyae and J. -C. Park, Phys. Lett. B **718**, 1425 (2013) [arXiv:1205.4151 [hep-ph]].
- [14] L. Wang and X. -F. Han, Phys. Rev. D **87**, 015015 (2013) [arXiv:1209.0376 [hep-ph]].
- [15] Rencontres de Moriond 2013 Electroweak Session,
<https://indico.in2p3.fr/conferenceDisplay.py?confId=7411>;
 Rencontres de Moriond 2013 QCD Session,
<http://moriond.in2p3.fr/QCD/2013/MorQCD13Prog.html>.
- [16] J. Ellis, T. You and , arXiv:1303.3879 [hep-ph].
- [17] M. Carena, I. Low and C. E. M. Wagner, “Implications of a Modified Higgs to Diphoton Decay Width,” arXiv:1206.1082 [hep-ph].
- [18] Z. Bern, P. Gondolo, M. Perelstein and , Phys. Lett. B **411**, 86 (1997) [hep-ph/9706538].
- [19] L. Bergstrom and P. Ullio, Nucl. Phys. B **504**, 27 (1997) [hep-ph/9706232].
- [20] J. Kumar, P. Sandick and , arXiv:1303.2384 [hep-ph].
- [21] T. Bringmann, L. Bergstrom and J. Edsjo, “New Gamma-Ray Contributions to Supersymmetric Dark Matter Annihilation,” JHEP **0801**, 049 (2008) [arXiv:0710.3169 [hep-ph]];
- L. Bergstrom, T. Bringmann, M. Eriksson and M. Gustafsson, “Gamma rays from heavy neutralino dark matter,” Phys. Rev. Lett. **95**, 241301 (2005) [hep-ph/0507229];
- A. Birkedal, K. T. Matchev, M. Perelstein and A. Spray, “Robust gamma ray signature of

- WIMP dark matter,” hep-ph/0507194.
- [22] B. Shakya, arXiv:1209.2427 [hep-ph].
 - [23] M. Carena, S. Gori, I. Low, N. R. Shah and C. E. M. Wagner, JHEP **1302**, 114 (2013) [arXiv:1211.6136 [hep-ph]].
 - [24] D. Das, U. Ellwanger and P. Mitropoulos, JCAP **1208**, 003 (2012) [arXiv:1206.2639 [hep-ph]].
 - [25] G. Chalons, M. J. Dolan and C. McCabe, JCAP **1302**, 016 (2013) [arXiv:1211.5154 [hep-ph]].
 - [26] U. Ellwanger, JHEP **1203**, 044 (2012) [arXiv:1112.3548 [hep-ph]];
K. Schmidt-Hoberg and F. Staub, JHEP **1210**, 195 (2012) [arXiv:1208.1683 [hep-ph]];
K. Choi, S. H. Im, K. S. Jeong and M. Yamaguchi, JHEP **1302**, 090 (2013) [arXiv:1211.0875 [hep-ph]].
 - [27] K. Schmidt-Hoberg, F. Staub and M. W. Winkler, JHEP **1301**, 124 (2013) [JHEP **1301**, 124 (2013)] [arXiv:1211.2835 [hep-ph]].
 - [28] S. Di Chiara and K. Hsieh, Phys. Rev. D **78**, 055016 (2008) [arXiv:0805.2623 [hep-ph]].
 - [29] A. Delgado, G. Nardini and M. Quiros, Phys. Rev. D **86**, 115010 (2012) [arXiv:1207.6596 [hep-ph]].
 - [30] Z. Kang, Y. Liu and G. -Z. Ning, arXiv:1301.2204 [hep-ph].
 - [31] L. Wang, X. -F. Han and , arXiv:1303.4490 [hep-ph].
 - [32] E. Aprile *et al.* [XENON100 Collaboration], Phys. Rev. Lett. **109**, 181301 (2012) [arXiv:1207.5988 [astro-ph.CO]].
 - [33] J. R. Espinosa and M. Quiros, Nucl. Phys. B **384**, 113 (1992).
 - [34] J. R. Espinosa and M. Quiros, Phys. Lett. B **279**, 92 (1992).
 - [35] A. Rajaraman, T. M. P. Tait and D. Whiteson, JCAP **1209**, 003 (2012) [arXiv:1205.4723 [hep-ph]].
 - [36] M. Su and D. P. Finkbeiner, arXiv:1206.1616 [astro-ph.HE].
 - [37] P. Ullio and L. Bergstrom, Phys. Rev. D **57**, 1962 (1998) [hep-ph/9707333].
 - [38] G. Belanger, F. Boudjema, P. Brun, A. Pukhov, S. Rosier-Lees, P. Salati, A. Semenov and , Comput. Phys. Commun. **182**, 842 (2011) [arXiv:1004.1092 [hep-ph]].
 - [39] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076 [astro-ph.CO].
 - [40] G. Hinshaw *et al.* [WMAP Collaboration], arXiv:1212.5226 [astro-ph.CO].
 - [41] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267**, 195 (1996) [hep-ph/9506380].
 - [42] G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005) [hep-ph/0404175].

- [43] J. R. Ellis, A. Ferstl and K. A. Olive, Phys. Lett. B 481, (2000) 304, [arXiv:hep-ph/0001005].
- [44] [ATLAS Collaboration], ATLAS-CONF-2012-109.
- [45] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Rev. Lett. **109**, 171803 (2012) [arXiv:1207.1898 [hep-ex]].
- [46] J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, “A Phenomenological Profile of the Higgs Boson,” Nucl. Phys. B **106** (1976) 292.
- [47] M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, “Low-Energy Theorems for Higgs Boson Couplings to Photons,” Sov. J. Nucl. Phys. **30** (1979) 711 [Yad. Fiz. **30** (1979) 1368].
- [48] A. Djouadi, “The Anatomy of electro-weak symmetry breaking. II. The Higgs bosons in the minimal supersymmetric model,” Phys. Rept. **459** (2008) 1 [hep-ph/0503173].